



**Republic of the Philippines  
DEPARTMENT OF HEALTH  
OFFICE OF THE SECRETARY**

**LIST OF EMERGENCY MEDICINES (Inpatient)**

**Medicines**

1) Ceftriaxone

- Inj.: 250 mg vial + 2 mL 1% solution of lidocaine (IM)  
(as disodium/sodium salt)
- 500 mg vial + 2 mL 1% solution of lidocaine (IM)  
(as disodium/sodium salt)
- 250 mg vial + 5 mL diluent (IV)  
(as disodium/sodium salt)
- 500 mg vial + 5 mL diluent (IV)  
(as disodium/sodium salt)
- 1 g vial + 3.5 mL 1% solution of lidocaine (IM)  
(as disodium/sodium salt)
- 1 g vial + 10 mL diluent (IV)  
(as disodium/sodium salt)

2) Cefuroxime

- Oral: 250 mg and 500 mg tablet (as axetil)
- 125 mg/5 mL granules for suspension,  
50 mL and 70 mL (as axetil)
- 125 mg granules for suspension, per sachet (as axetil)
- 250 mg/5 mL granules for suspension, 60 mL
- Inj.: 250 mg and 750 mg vial (IM, IV)
- 1.5 g vial (IV infusion)

3) Azithromycin

- Oral: 250 mg capsule (as base\*/as dihydrate)
- 500 mg tablet (as base\*/as dihydrate)
- 200 mg/5 mL powder for suspension, 15 mL and 30 mL (as base\*/as dihydrate)
- Inj.: 500 mg powder, vial (IV infusion) (as base\*/as dihydrate)

4) Doxycycline

- Oral: 50 mg and 100 mg capsule (as hyclate)

5) Amikacin

- Inj.: 50 mg/mL, 100 mg/mL, 125 mg/mL and 250 mg/mL, 2 mL ampul/vial (IM, IV) (as sulfate)
- 100 mg, 250 mg and 500 mg vial (IM, IV) (as sulfate)

6) Amoxicillin 500 mg capsule – approve (also include suspension for pedia)

- Oral: 250 mg and 500 mg tablet  
 100 mg/mL granules/powder for drops (suspension), 10 mL and 15 mL (as trihydrate)  
 125 mg/5 mL granules/powder for suspension, 30 mL and 60 mL (as trihydrate)  
 250 mg/5 mL granules/powder for suspension, 30 mL and 60 mL (as trihydrate)
- 7) Ampicillin 500 mg + Sulbactam  
 Inj.: 250 mg ampicillin + 125 mg sulbactam (as sodium salt) per vial (IM, IV)  
 500 mg ampicillin + 250 mg sulbactam (as sodium salt) per vial (IM, IV)
- 8) Co-amoxiclav  
 Oral: 500 mg Amoxicillin (as trihydrate) + 125 mg potassium clavulanate per tablet  
 875 mg amoxicillin (as trihydrate) + 125 mg potassium clavulanate per tablet  
 125 mg amoxicillin (as trihydrate) + 31 mg potassium clavulanate per 5 mL granules/powder for suspension, 30 mL and 60 mL  
 200 mg amoxicillin (as trihydrate) + 28.5 mg potassium clavulanate per 5 mL granules/powder for suspension, 70 mL  
 250 mg amoxicillin (as trihydrate) + 62.5 mg potassium clavulanate per 5 mL granules/powder for suspension, 60 mL and 100 mL  
 400 mg amoxicillin (as trihydrate) + 57 mg potassium clavulanate per 5 mL granules/powder for suspension, 30 mL and 70 mL
- 9) Oxacillin  
 Inj.: 250 mg and 500 mg vial (IM, IV) (as sodium salt)
- 10) Cloxacillin  
 Oral: 250 mg and 500 mg capsule (as sodium salt)  
 125 mg /5 mL powder for syrup/suspension, 60 mL (as sodium salt)
- 11) Benzyl penicillin  
 Inj.: 500, 000 units via (IM, IV) (as sodium salt)  
 1, 000, 000 units vial (IM, IV) (as sodium salt)  
 5, 000, 000 units vial (IM, IV) (as sodium salt)
- 12) Metronidazole  
 Oral: 250 mg and 500 mg tablet  
 125 mg/ base/5 mL (200 mg/5 mL) (as benzoate) suspension, 30 mL and 60 mL  
 Inj.: 5 mg/mL, 100 mL vial (IV infusion)
- 13) Clindamycin  
 Oral: 150 mg and 300 mg capsule (as hydrochloride)

75 mg / 5 mL granules for suspension, 60 mL (as palmitate hydrochloride)  
Inj.: 150 mg/mL, 2 mL ampul/vial and 4 mL ampul (IM, IV) (as phosphate)

14) Calcium gluconate

Inj.: 10%, 10 mL ampul/vial (IV)  
10%, 20 mL and 25 mL bottle (IV)

15) Sodium bicarbonate

Inj.: 1 mEq/mL, 10 mL vial (pediatric), 50 mL and 100 mL ampul/vial (adult) (IV infusion)

16) Hydrocortisone

Inj.: 50 mg/mL, 2 mL vial (IM, IV)(as sodium succinate)  
125 mg/mL, 2 mL and 4 mL vial (IV) (as sodium succinate)  
powder, 100 mg, 250 mg and 500 mg vial (IV) (as sodium succinate)  
Cream or Ointment: 1%, 5g and 10 g tube and 500 g jar

17) Furosemide

Oral: 20 mg and 40 mg tablet  
Inj.: 10 mg/mL, 2 mL ampul (IM, IV)

18) Ketorolac

Inj.: 30 mg/mL, 1 mL ampul (IM, IV) (as tromethamol)

19) Phenytoin

Oral: 30 mg and 100 mg capsule (as sodium salt)  
30 mg/5 mL suspension, 120 mL (as sodium salt)  
Inj.: 50 mg/mL, 2 mL ampul (IV) (as sodium salt)

20) Phytomenadione

Oral/Inj.: 2mg/0.2 mL pediatric ampul (IM, IV, PO) (as mixed micelle)  
Inj.: 10 mg/mL, 1 mL ampul (IM, IV) (as aqueous colloidal solution with benzyl alcohol)  
10 mg/mL, 1 mL ampul (IM, IV) (as mixed micelle)

21) Potassium chloride

Inj.: 2 mEq/mL, 2 mL and 5 mL ampul (IV infusion)  
2 mEq/mL, 20 mL and 10 mL vial (IV infusion)

22) Tranexamic acid

Oral: 250 mg and 500 mg capsule  
500 mg tablet  
Inj.: 100 mg/mL, 2.5 mL and 5 mL ampul (IM, IV)

23) D50 water 50 cc.

24) Sterile Water for Injection, 50 mL

25) NSS

250 mL, 500 mL and 1 L bottle/bag (IV infusion)

D5LR

5% Dextrose In Lactated Ringers

Inj.: 250 mL, 500 mL and 1 L bottle/bag (IV infusion)

LR

250 mL, 500 mL and 1 L bottle/bag (IV infusion)

D5NSS

250 mL, 500 mL and 1 L bottle/bag (IV infusion)

D5W

250 mL, 500 mL and 1 L bottle/bag (IV infusion and as vehicle for IV medications)

26) Dopamine

Inj.: 40 mg/mL, 5 mL vial/ampul (IV) (as hydrochloride)

80 mg/mL, 5 mL vial (IV) (as hydrochloride)

800 micrograms/mL, 250 mL D5W (pre-mixed) (IV) (as hydrochloride)

1.6 mg/mL, 250 mL D5W (pre-mixed) (IV) (as hydrochloride)

3.2 mg/mL, 250 mL D5W (pre-mixed) (IV) (as hydrochloride)

27) Dobutamine

Inj.: 12.5mg/mL, 20mL vial (IV) (as hydrochloride)

50mg/mL, 5mL ampul, 5 mL ampul (concentrate) (IV infusion) (as hydrochloride)

1 mg/mL, 250 mL D5W (pre-mixed) (IV) (as hydrochloride)

2 mg/mL, 250 mL D5W (pre-mixed) (IV) (as hydrochloride)

4 mg/mL, 250 mL D5W (pre-mixed) (IV) (as hydrochloride)

28) Nicardipine

Oral: 10 mg and 20 mg tablet (as hydrochloride)

Inj.: 1mg/mL, 2 mL and 10mL ampul (IV) (as hydrochloride)

29) Epinephrine

Inj.: 1 mg/mL, 1 mL ampul (IM, SC) (as hydrochloride)

30) Isosorbide dinitrate

Oral: 5mg, 10 mg and 20 mg tablet

20 mg and 40 mg MR tablet/capsule

Sublingual: 5 mg tablet

Inj.: 1 mg/mL, 10 mL ampul (IV)

31) Salbutamol

Oral: 2 mg tablet (as sulfate)

2 mg/5 mL syrup, 30 mL, 60 mL and 120 mL (as sulfate)

Inhalation:

- Dry Powder Inhaler (DPI):  
200 micrograms/dose (as sulfate) with appropriate accompanying dispenser  
400 micrograms/dose (as sulfate) with appropriate accompanying dispenser
- Metered Dose Inhaler (MDI):  
100 micrograms/dose x 200 and 300 doses (as sulfate)
- Breath Actuated MDI:  
100 micrograms/dose x 200 doses (as sulfate)  
100 micrograms/dose x 400 doses (as sulfate)
- Resp. Soln.: (for nebulization)  
1 mg/mL, 2.5 mL (unit dose) (as sulfate)  
1 mg/mL, 30 mL bottle (as sulfate)  
2 mg/mL, 2.5 mL (unit dose) (as sulfate)  
5 mg/mL, 10 mL and 20 mL (multidose) (as sulfate)

32) Salbutamol/Ipratropium

MDI: 21 micrograms ipratropium (as bromide) + 120 micrograms salbutamol x 200 doses x 10 mL

Resp. Soln.: (for nebulization)

500 micrograms ipratropium (as bromide anhydrous) +  
2.5 mg salbutamol (as base) x 2.5 mL (unit dose)

33) Betahistine

Oral: 6 mg tablet (as mesilate)

8 mg and 24 mg tablet (as hydrochloride)

34) Metoclopramide

Oral: 10 mg tablet (as hydrochloride)

5 mg/5 mL syrup, 60 mL (as base and as hydrochloride)

Inj.: 5 mg/mL, 2 mL and 3 mL ampul (IM, IV) (as base and as hydrochloride)

35) Omeprazole

Oral: 10 mg, 20 mg and 40 mg capsule

Inj.: powder, 40 mg vial + 10 mL solvent ampul (IV)

36) Hyoscine

Oral: 10 mg tablet (as N-butyl bromide)

5 mg/5 mL syrup, 60 mL (as N-butyl bromide)

Inj.: 20 mg/mL, 1 mL ampul (IM, IV, SC) (as N-butyl bromide)

500 micrograms/mL, 1 mL ampul (IM, IV) (as hydrobromide)

200 micrograms/mL, 1 mL ampul (IM, IV) (as hydrobromide)

37) Mefenamic acid

Oral: 250 mg and 500 mg tablet/capsule

38) Ibuprofen

Oral: 200 mg and 400 mg tablet  
800 mg MR table  
100 mg/5 mL suspension , 60 mL

39) Tramadol

Oral: 50 mg capsule (as hydrochloride)  
100 mg, 150 mg and 200 mg MR tablet (as hydrochloride)  
Inj.: 50 mg/mL, 1 mL and 2 mL ampul (IM, IV, SC) (as hydrochloride)

40) Paracetamol

Oral: 300 mg and 500 mg tablet  
100 mg/mL drops, 15 mL (alcohol-free)  
120 mg/5 mL (125 mg/5 mL) syrup/suspension, 30 mL, 60 mL and 120 mL  
(alcohol-free)  
250 mg/5 mL syrup/suspension, 30 mL, 60 mL and 120 mL (alcohol-free)

41) Diphenhydramine

Oral: 25 mg and 50 mg capsule (as hydrochloride)  
12.5 mg/5 mL, 60 mL and 120 mL syrup  
Inj.: 50 mg/mL, 1 mL ampul (IM, IV) (as hydrochloride)

42) Chlorpromazine

Oral: 50 mg, 100 mg and 200 mg tablet (as hydrochloride)  
Inj.: 25 mg/mL, 1 mL ampul (IM, IV) (as hydrochloride)

43) Diazepam

Oral: 2 mg, 5 mg and 10 mg tablet  
Inj.: 5 mg/mL, 2 mL ampul (IM, IV)

44) Valproic acid

Oral: 250 mg (as disodium salt or as valproic acid)  
250 mg/ 5 mL syrup, 120 mL (as valproic acid)

45) Mannitol

Inj.: 20%, 250 mL and 500 mL bottle (IV)

46) Propranolol

Oral: 10 mg, 40 mg and 80 mg tablet (as hydrochloride)

40 mg mg tablet and 80 mg MR capsule (as hydrochloride)

47) Digitalis

Oral: 250 microgram tablet

50 micrograms/mL elixir, 60 mL

Inj.: 250 micrograms/mL, 2 mL ampul (IM, IV)

Aspirin

Oral: 80 mg tablet

49) Clopidogrel

Oral: 75 mg tablet

50) Atropine

Inj.: 500 micrograms/mL and 1 mg / mL, 1 mL ampul (IM, IV) (as sulfate)

51) Oxytocin (*synthetic*)

Inj.: 5 IU / mL and 10 IU / mL, 1 mL ampul (IM, IV)

52) Magnesium sulphate

Inj.: 250 mg/mL, 2 mL , 10 mL ampul and 20 mL vial (as heptahydrate)

500 mg/mL, 2 mL and 10 mL ampul (as heptahydrate)

53) Insulin - approve

**Short Acting**

Regular, Insulin (recombinant DNA, human)

100 IU/mL 3 mL pre-filled syringe (IM,IV, SC)

100 IU/mL, 10 mL vial (IM,IV, SC)

**Intermediate Acting (recombinant DNA)**

Isophane Insulin Human (recombinant DNA)

100 IU/mL, 3 mL pre-filled syringe (IM, SC)

100 IU/mL, 10 mL vial (IM, SC)

Biphasic isophane human insulin 70/30 (recombinant DNA)

70% isophane suspension + 30% soluble insulin in

100 IU/mL vial (IM, SC) and 100 IU/mL, 3 mL disposable syringe (IM,SC)

Insulin zinc suspension human

100 IU /mL, 10 mL vial (IM, SC)

54) Oral rehydration solutions

Composition of reduced osmolarity ORS per liter of water (WHO recommended)

Sodium chloride – 2.6 g

Trisodium citrate dehydrate – 2.9 g

Potassium chloride – 1.5 g

Glucose anhydrous – 13.5 g

**Total Weight – 20.5 g**

Reduced osmolarity ORS

Equivalent in mmol/L:  
Sodium – 75  
Chloride – 65  
Potassium – 20  
Citrate – 10  
Glucose anhydrous – 75  
**Total osmolarity – 245**

55) Silver sulfadiazine

Cream: 1%, 5 g, 10 g, 15 g, 20 g, 25 g, 30 g and 50 g tube  
400 g, 450 g and 500 g jar (micronized)

56) Tetanus Immunoglobulin (Human)

Inj.: 1000 IU/ mL, 1.5 mL vial (IM)  
1500 IU/mL, 1 mL ampul (IM)  
250 IU/mL, 1 mL, 2 mL and 4 mL ampul (IM)  
250 IU/mL, 1 mL pre-filled syringe (IM)  
250 units/mL, 1 mL and 2 mL vial (IM)

57) Tetanus Toxoid

Inj.: 0.5 mL ampul (IM)  
5 mL and 10 mL vial

58) Povidone Iodine

Solution: 10%, 15 mL, 30 mL, 60 mL, 120 mL, 1 L and 1 gallon bottle  
Surgical Cleanser: 7.5%, 60 mL, 120 mL, 480 mL, 1 L and 1 gallon bottle  
(Note: Povidone Iodine Swabs are preferred)

59) Clonidine

Oral: 75 microgram and 150 microgram tablet (as hydrochloride)  
Inj.: 150 micrograms/mL, 1 mL ampul (IV) (as hydrochloride)

60) Captopril

Oral: 25 mg and 50 mg tablet

61) Amlodipine

Oral: 5mg and 10 mg tablet (as besilate/camsylate)

62) Losartan

Oral: 50 mg and 100 mg tablet (as potassium salt)

**Local Anesthetics**

1) Bupivacaine

Inj.: 0.5%, 5 mL and 10 mL ampul (local infiltration) (as hydrochloride)  
0.5%, 5 mL, 10 mL, 20 mL and 50 mL ampul/vial (local infiltration) (as hydrochloride)  
0.5% 4 mL ampul (spinal) with 8% dextrose (as hydrochloride)



0.5% (isobaric), 5 mL ampul (spinal) (as hydrochloride)

2) Lidocaine

Inj.: 1%, 5 mL ampul and 50 mL vial (local infiltration) (as hydrochloride)

2%, 5 mL, 10 mL and 50 mL vial (epidural, local infiltration) (as hydrochloride)

2%, 1.8 mL carpule (with epinephrine) (local infiltration)

Jelly: 2%, 30g (as hydrochloride)

Ointment: 5%, 35 g and 50 g (as hydrochloride)

Solution: 4%, 30 mL (topical solution) (as hydrochloride)

Spray: 10%, 50 mL (as hydrochloride)

3) Tetracaine

Inj.: 20 mg ampul (sterile powder) (spinal) (as hydrochloride)

**Peritoneal Dialysis Apparatus (Solutions and Equipment)**

A. Peritoneal Access Devices:

1. Acute catheters
2. Chronic catheters (usually coiled type)

B. Dialysis Solutions: May keep at room temperature but must be warmed to body temperature before instilling

1. 1.5% (2 liter bag)
2. 2.3% (2 liter bag)
3. 4.25% (2 liter bag)

C. Transfer Sets:

1. PD straight extension catheter (40 cm)
2. Y set extension tubing

D. Connectors and Disinfection Caps

E. PD Organizer

F. Weighing Scale (5 kg) - small hand-held type

G. Plastic Basin - approx. 1.5 feet width, for holding dialysis bags

H. Others:

1. OR pack
2. Dressing kit - cotton applicator, gauze 4x4, transparent dressing g10x25 cm), Gauze 4x4, micro pore tape
3. Sterile gloves
4. Surgical mask
5. 1 cc syringes with needle (for injecting heparin)
6. Heating pad for warming the dialysis solutions

I. Medicines:

1. Heparin
2. Tranexamic acid
3. Antibiotics when necessary: Amikacin, cefuroxime
4. Betadine solution

## **Medical Supplies and Equipments**

- 1) Syringe 2,5,10cc
- 2) Tuberculin syringe
- 3) Insulin syringe (0.3, 0.5, 1 cc)
- 4) CBG strips (should be compatible with glucometer)
- 5) IV Cannula
- 6) Micropore
- 7) Soluset
- 8) Sutures
- 9) OR Packs
- 10) Gloves
- 11) Gauze
- 12) Bandages
- 13) Nebulizing Kit
- 14) Oxygen Tank
- 15) Glucometer
- 16) Urinary Catheters
- 17) Nasogastric Tubes (NGTs)
- 18) Nasal cannula
- 19) Face mask for O<sub>2</sub> (adult and pedia)
- 20) Chest tube insertions
- 21) PD Catheter
- 22) Cotton
- 23) Eye Patch/Eye Shield
- 24) Splints
- 25) Tongue Depressor
- 26) External Fixators
- 27) Canes/ Crutches/ Walkers

### **Accessories:**

- 1) Battery
- 2) Candle
- 3) Flashlight
- 4) Rechargeable Lamp
- 5) Matches
- 6) Solar powered lamps