



**Republic of the Philippines  
DEPARTMENT OF HEALTH  
OFFICE OF THE SECRETARY**

**LIST OF EMERGENCY MEDICINES (Outpatient)**

**A. Wounds/injuries**

- 1) Povidone iodine  
Solution: 10%, 15 mL, 30 mL, 60 mL, 120 mL, 240 mL, 1 L and 1 gallon bottle  
Surgical Skin Cleanser:  
7.5%, 60 mL, 120 mL, 480 mL, 1 L and 1 gallon bottle
- 2) Lidocaine  
Inj.: 1%, 5 mL and 20 mL ampul (as hydrochloride)  
2%, 2 mL, 5 mL and 20 mL ampul (as hydrochloride)  
2%, 10 mL, 20 mL and 50 mL vial (as hydrochloride)
- 3) Ethyl Alcohol solution  
95% for dilution to 70% (with BIR seal)
- 4) Cloxacillin  
Oral: 250 mg and 500 mg capsule (as sodium salt)  
125 mg/5 mL powder for syrup/suspension, 60 mL (as sodium salt)
- 5) Tetanus toxoid  
Inj.: 0.5 mL ampul (IM)  
5 mL and 10 mL vial (IM)
- 6) Anti-Tetanus Serum (Equine)  
Inj.: 4000 IU/mL, 2.5 mL vial (IM)  
1500 IU/mL, 1 mL and 1.5 mL vial/ampul (IM)
- 7) Hydrogen peroxide  
Solution: 3%, 60 mL and 120 mL bottle
- 8) Cobra antivenin  
Inj.: 800 MU/ 4.8 mL, 1 mL ampul (for IV infusion)
- 9) 0.9% Sodium Chloride (Normal Saline Solution)  
Inj.: 2 mL, 2.5 mL, 5 mL, 10 mL and 20 mL ampul  
50 mL, 100 mL, 200 mL, 250 mL, 500 mL and 1 L bottle/bag (IV infusion)  
Composition:  
Na<sup>+</sup> --- 154 mmol/L  
Cl<sup>-</sup> --- 154 mmol/L
- 10) Silver sulfadiazine  
Cream: 1%, 5 g, 10 g, 15 g, 20 g, 25 g, 30 g and 50 g tube  
400 g, 450 g and 500 g jar (micronized)

11) Mupirocin

Cream: 2%, 5 g sachet and 15 g tube

Ointment: 2%, 5 g and 15 g tube

**B. Diarrhea**

1) Oral Rehydration Salts (ORS 75-replacement)

2) Metronidazole

Oral: 250 mg and 500 mg tablet, 125 mg base/5 mL (200 mg/5 mL (as benzoate) suspension, 30 mL and 60 mL

Inj.: 5 mg/mL, 100 mL vial (IV infusion)

Rectal: 1 g suppository

3) Zinc sulfate tablet and syrup

Oral: tablet, (equiv. to 30 mg elemental zinc) (as gluconatetrihydrate)

solution, (equiv. to 10 mg elemental zinc/mL) drops, 15 mL (as sulfate monohydrate)

solution, (equiv. to 20 mg elemental zinc/5 mL) syrup, 60 mL (as sulfate monohydrate)

4) Ciprofloxacin (for adult use)

Oral: 250 mg and 500 mg tablet (as hydrochloride)

Inj.: 2 mg/mL, 50 mL and 100 mL vial (IV infusion) (as lactate)

5) Loperamide (for adult use)

Oral: 2 mg capsule (as hydrochloride) (N.B.: Not for infants and children less than 12 years old)

6) Tetracycline

Oral: 250 mg and 500 mg capsule

7) Sodium dichloro-isocyanurate (for water purification)

Solution: 3.5 mg tablet (free available isocyanurate chlorine 2 mg)

8.68 mg tablet (free available chlorine 5 mg)

12.5 mg tablet (free available chlorine 8 mg)

17 mg tablet (free available chlorine 10 mg)

67 mg tablet (free available chlorine 40 mg)

**C. URTI/Pneumonia**

1) Amoxicillin

250 mg and 500 mg capsule (as trihydrate)

100 mg/mL granules/powder for drops (suspension), 10 mL and 15 mL (as trihydrate)

125 mg/5 mL granules/powder for suspension, 30 mL and 60 mL (as trihydrate)

250 mg/5 mL granules/powder for suspension, 30 mL and 60 mL (as trihydrate)

2) Cefuroxime

250 mg and 500 mg tablet (as axetil)

125 mg/5 mL granules for suspension,

50 mL and 70 mL (as axetil)  
125 mg granules for suspension, per sachet (as axetil)  
250/5 mL granules for suspension, 60 mL  
Inj.: 250 mg and 750 mg vial (IM, IV)  
1.5 g vial (IV infusion) (as sodium salt)

- 3) Co-amoxiclav (amoxiclav + potassium clavulanate)  
500 mg amoxicillin (as trihydrate) + (125 mg potassium clavulanate potassium per tablet)  
875 mg amoxicillin (as trihydrate) + 125 mg potassium clavulanate per tablet  
125 mg amoxicillin (as trihydrate) + 31 mg potassium clavulanate, per 5 mL granules/powder  
for suspension, 30 mL and 60 mL  
200 mg amoxicillin (as trihydrate) + 28.5 mg potassium clavulanate per 5 mL  
granules/powder for suspension, 70 mL  
250 amoxicillin (as trihydrate) + 62.5 mg potassium clavulanate per 5 mL granules/powder  
for suspension, 60 mL and 100 mL  
400 amoxicillin (as trihydrate) + 57g potassium clavulanate per 5 mL granules/powder  
for suspension, 30 mL and 70 mL
- 4) Cotrimoxazole (*sulfamethoxazole + trimethoprim*)  
(also for *Pneumocystis jiroveci (carinii) and chlamydiae*)  
400 mg sulfamethoxazole + 80 mg trimethoprim per tablet/capsule  
800 mg sulfamethoxazole + 160 mg trimethoprim per tablet  
200 mg sulfamethoxazole + 40 mg trimethoprim/5 mL suspension, 30ml, 60ml and 100m  
400 mg sulfamethoxazole + 80 mg trimethoprim/5 mL suspension, 30 mL and 60 mL  
Inj.: 400 mg sulfamethoxazole + 80 mg trimethoprim, 5 mL ampul (IV infusion)

#### **D. Other common infections**

- 1.) Doxycycline (also for cholera and leptospirosis)  
50 mg and 100 mg capsule (as hyclate)
- 2.) Clarithromycin  
Oral: 250 mg and 500 mg base tablet  
500 mg MR tablet  
125 mg/5 mL granules/powder for suspension, 25 mL, 50 mL and 70 mL
- 3.) Ciprofloxacin  
250mg and 500 mg tablet (as hydrochloride)  
Inj.: 2 mg/mL, 50 mL and 100 mL vial (IV infusion) (as lactate)

#### **E. Diabetes**

- 1) Metformin  
500 mg tablet/film coated tablet (as hydrochloride)  
850 mg and 1 g tablet (as hydrochloride)
- 2) Gliclazide  
30 mg MR tablet  
80 mg tablet

- 3) Insulin, Regular (recombinant DNA, human)  
Inj.: 100 IU/mL, 3 mL pre-filled syringe (IM, SC, IV)  
100 IU/mL, 10 mL vial (IM, SC, IV)  
*\*ensure cold chain*

## **F. Hypertension**

- 1) Amlodipine  
5 mg and 10 mg tablet (as besilate/camsylate)
- 2) Metoprolol  
50 mg tablet and 100 mg tablet (as tartrate)
- 3) Enalapril  
5 mg and 10 mg tablet and 20 mg tablet (as maleate)
- 4) Hydrochlorothiazide  
25 mg and 50 mg tablet
- 5) Clonidine  
75 microgram and 150 microgram tablet (as hydrochloride)  
Inj.: 150 micrograms/mL, 1 mL ampul (IV) (as hydrochloride)
- 6) Captopril (for hypertensive emergency)  
Oral: 25 mg and 50 mg tablet

## **G. Coronary Heart Disease**

- 1) Isosorbide dinitrate  
5 mg, 10 mg and 20 mg tablet  
20 mg and 40 mg MR tablet/capsule  
Sublingual: 5 mg tablet  
Inj.: 1 mg/mL, 10 mL ampul (IV)
- 2) Aspirin  
Oral: 80 and 100 mg enteric coated tablet, 300 mg tablet
- 3) Simvastatin  
Oral: 10 mg, 20 mg and 40 mg tablet

## **H. Asthma**

- 1) Salbutamol  
2 mg tablet (as sulfate)  
2 mg/5 mL syrup, 30 mL, 60 mL and 120 mL (as sulfate)  
  
Inhalation:  
Dry Powder Inhaler (DPI):  
200 micrograms/dose (as sulfate) with appropriate accompanying dispenser  
400 micrograms/dose (as sulfate) with appropriate accompanying dispenser  
Metered Dose Inhaler (MDI):

100 micrograms/dose x 200 and 300 doses (as sulfate)

**Breath Actuated MDI:**

100 micrograms/dose x 200 doses (as sulfate)

100 micrograms/dose x 400 doses (as sulfate)

**Resp. Soln.: (for nebulization)**

1 mg/mL, 2.5 mL (unit dose) (as sulfate)

1 mg/mL, 30 mL bottle (as sulfate)

2 mg/mL, 2.5 mL (unit dose) (as sulfate)

5 mg/mL, 10 mL and 20 mL (multidose) (as sulfate)

**2) Fluticasone + salmeterol**

**Inhalation:**

**Dry Powder Inhaler (DPI):**

100 micrograms fluticasone (as propionate) + 50 micrograms salmeterol (as xinafoate) x 28 doses and 60 doses with appropriate accompanying dispenser

250 micrograms fluticasone (as propionate) + 50 micrograms salmeterol (as xinafoate) x 28 doses and 60 doses with appropriate accompanying dispenser

500 micrograms fluticasone (as propionate) + 50 micrograms salmeterol (as xinafoate) x 28 doses and 60 doses with appropriate accompanying dispenser

**Metered Dose Inhaler (MDI):**

50 micrograms fluticasone (as propionate) + 25 micrograms salmeterol (as xinafoate) x 120 doses

125 micrograms fluticasone (as propionate) + 25 micrograms salmeterol (as xinafoate) x 120 doses

250 micrograms fluticasone (as propionate) + 25 micrograms salmeterol (as xinafoate) x 120

**I. Epilepsy / Convulsion**

**1) Diazepam**

Inj.: 5 mg/mL, 2 mL ampul (IV)

Rectal: 5 mg/2.5 mL and 10 mg/2.5 mL in rectal tube

**J. Pain/fever**

**1) Paracetamol**

300 mg (325 mg) and 500 mg tablet

120 mg (125 mg)/5 mL syrup/suspension, 60 mL (alcohol-free)

250 mg/5 mL syrup/suspension, 60 mL (alcohol-free preferred)

100 mg/mL drops, 15 mL (alcohol-free)

Rectal: 125 mg & 250 mg suppository

**2) Mefenamic acid**

250 mg and 500 mg tablet/capsule

- 3) Ibuprofen  
200 mg and 400 mg tablet  
100 mg/5 mL suspension, 60 mL
- 4) Celecoxib  
Oral: 100 mg, 200 mg and 400 mg capsule/tablet

#### **K. Acid Peptic Disease**

- 1) Omeprazole  
Oral: 10 mg, 20 mg and 40 mg capsule  
Inj.: powder, 40 mg vial + 10 mL solvent ampul (IV)
- 2) Magnesium aluminum hydroxide  
Oral: 225 mg aluminum hydroxide + 200 mg magnesium hydroxide, per 5 mL suspension, 60 mL, 120 mL, 180 mL, 250 mL, 355 mL

#### **L. Anti Allergy**

- 1) Diphenhydramine  
Oral: 25 mg and 50 mg capsule (as hydrochloride)  
12.5 mg/5 mL syrup, 30 mL, 60 mL and 120 mL (alcohol-free) (as hydrochloride)  
Inj.: 50 mg/mL, 1 mL ampul and 10 mL vial (IM, IV) (as hydrochloride)
- 2) Loratadine  
Oral: 10 mg tablet and 10 mg film coated tablet  
5 mg/5 mL syrup, 30 mL
- 3) Cetirizine  
Oral: 10 mg tablet (as dihydrochloride)  
10 mg/mL drops, 10 mL (as dihydrochloride)  
1 mg/mL solution, 30 mL and 60 mL (as dihydrochloride)  
5 mg/5 mL syrup, 30 mL (as dihydrochloride)
- 4) Epinephrine  
Inj.: 1 mg/mL, 1 mL ampul (IM, SC) (as hydrochloride)  
0.3 mg auto-injector (IM-Preloaded)

#### **M. Psychotropics**

- 1) Diazepam  
Oral: 2 mg, 5 mg and 10 mg tablet  
Inj.: 5 mg/mL, 2 mL ampul (IM, IV)
- 2) Hydroxyzine  
Oral: 10 mg and 25 mg tablet (as dihydrochloride)  
2 mg/mL syrup, 60 mL (as dihydrochloride or as hydrochloride)  
Inj.: 50 mg/mL, 1 mL vial (IM, IV)
- 3) Olanzapine

Oral: 5 mg and 10 mg tablet  
Inj.: 10 mg vial (IM)

#### **N. Ophthalmic**

- 1) Fusidic acid  
Eye Drops Suspension:  
1%, 5 g tube (as sulfate)
- 2) 0.9% Sodium Chloride (Normal Saline Solution)  
Inj.: 2 mL, 2.5 mL, 5 mL, 10 mL and 20 mL ampul  
50 mL, 100 mL, 200 mL, 250 mL, 500 mL and 1 L bottle/bag (IV infusion)  
Composition:  
Na<sup>+</sup> --- 154 mmol/L  
Cl<sup>-</sup> --- 154 mmol/L

#### **O. Otic**

- 1) Neomycin + Polymyxin B + Fluocinolone acetonide  
Ear Drops Solution:  
3.5 mg neomycin (as sulfate) + 10,000 units polymyxin B (as sulfate) + 0.025% fluocinolone acetonide/mL, 5 mL bottle

#### **P. Antifungal**

- 1) Clotrimazole  
Cream: 1%-2%, 3.5 g, 15 g and 450 g tube  
Lotion: 10 mL  
Shampoo: 6 mL and 10 mL sachet, 60 mL and 100 mL bottle  
Solution: 10 mg/mL, 15 mL and 25 mL bottle  
Vaginal: 100 mg, 300 mg and 500 mg ovule
- 2) Sulfur ointment  
Cream or Ointment: 5%, 15 g and 30 g tube

#### **Q. Scabicides**

- 1) Permethrin  
Lotion: 1%, 125 mL bottle  
5%, 30 mL and 60 mL bottle  
Shampoo (Crema Rinse): 1%, 30 mL and 60 mL bottle

#### **R. Oral contraceptives**

- 1) ETHINYLESTRADIOL +LEVONORGESTREL  
Oral: 30 microgram ethinylestradiol + 150 microgram levonorgestrel per tablet